arXiv:1103.2051v1 [math.GR] 10 Mar 2011

Regular tessellations of the hyperbolic plane by fundamental domains of a Fuchsian group

Robert Yuncken
Penn State University

January 14, 2013

Running Head: “Regular tessellations and Fuchsian groups”
Keywords: FUCHSIAN GROUP, REGULAR TESSELLATION, HYPERBOLIC PLANE, FUNDAMENTAL DOMAIN.
AMS Subject Classification: 20H10—Fuchsian groups and their generalizations.

Abstract

For positive integers p and q with $1/p + 1/q < 1/2$, a tessellation of type $\{p, q\}$ is a tessellation of the hyperbolic plane by regular p-gons with q p-gons meeting at each vertex. In this paper, a necessary and sufficient condition on the integers p and q is established to determine when a tessellation of type $\{p, q\}$ can be realized as a tessellation of the hyperbolic plane by fundamental domains of some Fuchsian group. Specifically, a tessellation of type $\{p, q\}$ is a tessellation by fundamental domains if and only if q has a prime divisor less than or equal to p.

It is well-known that, for integers p and q with $1/p + 1/q < 1/2$, the hyperbolic plane $\mathbb{H}^2$ can be tessellated by regular p-gons with q p-gons meeting at each vertex. This is called a tessellation of type $\{p, q\}$. The group of all orientation-preserving symmetries of a tessellation of type $\{p, q\}$ is a Fuchsian group, and a tessellation of $\mathbb{H}^2$ by fundamental domains for this group is obtained by subdividing each of the regular p-gons in the original tessellation into p triangles sharing a vertex at its centre.

The question we deal with here is whether the p-gons themselves are fundamental domains for some Fuchsian group. Equivalently, one can ask for which p and q does there exist a subgroup of the full group of symmetries of the tessellation of type $\{p, q\}$ which acts freely and transitively on the set of p-gons in the tessellation? The following result answers this question.

Theorem 0.1. *The tessellation of type $\{p, q\}$ is a tessellation of the hyperbolic plane by fundamental domains of some Fuchsian group if and only if q has a prime divisor less than or equal to p.*

The proof of this theorem can be broken into two Lemmas.

Lemma 0.2. *The tessellation of type $\{p, q\}$ is a tessellation of the hyperbolic plane by fundamental domains of some Fuchsian group if and only if there exists an involution σ in S_p (the symmetric group on p letters) such that $(\sigma\rho)^q = 1$, where ρ is the cyclic permutation $(1\ 2\ \dots\ p)$ of order p.*

Proof. Let us fix a p-gon $\mathcal{F}$ in the tessellation. Let $v_1, \ldots, v_p$ be the vertices of $\mathcal{F}$, labeled clockwise from some arbitrary vertex v_1. Let e_1 denote the edge between v_p and v_1, and let e_i denote the edge between v_{i-1} and v_i for $i = 2, \ldots, p$.

First, let us prove the necessity of the combinatorial condition. Suppose that $\mathcal{F}$ is a fundamental domain for some Fuchsian group Γ. For each of the p p-gons adjacent to $\mathcal{F}$, there is a unique element of Γ which maps it onto $\mathcal{F}$. This induces the well-known edge-pairing for a (polygonal) fundamental domain—namely, for each edge e_i of $\mathcal{F}$ there is a unique non-identity element $\gamma_i \in \Gamma$ and a unique edge of $\mathcal{F}$, which we will denote by $e_{\sigma(i)}$, such that γ_i maps $e_{\sigma(i)}$ onto e_i. Note that we do not exclude the possibility that $e_i = e_{\sigma(i)}$, in which case γ_i is a rotation about the midpoint of e_i by π.

Since γ_i^{-1} maps e_i onto $e_{\sigma(i)}$, we see that $\gamma_i^{-1} = \gamma_{\sigma(i)}$ and $e_{\sigma(\sigma(i))} = e_i$. Thus the edge-pairing of $\mathcal{F}$ provides an involution $\sigma \in S_p$. We now prove the relation between σ and ρ by considering the q images of $\mathcal{F}$ about any vertex v_i.

Consider any edge e_i of $\mathcal{F}$. The adjacent edge of $\mathcal{F}$ which shares the vertex v_i is $e_{\rho(i)}$, where ρ is the cyclic permutation defined in the statement of the theorem. By the definition of σ, $\gamma_{\sigma\rho(i)}\mathcal{F}$ is the p-gon adjacent to $\mathcal{F}$ in the tessellation which shares the edge $e_{\rho(i)}$, and $\gamma_{\sigma\rho(i)} v_{\sigma\rho(i)} = v_i$. The edge of $\gamma_{\sigma\rho(i)}\mathcal{F}$ which is adjacent to $\gamma_{\sigma\rho(i)} e_{\sigma\rho(i)}$ and shares the vertex v_i is $\gamma_{\sigma\rho(i)} e_{\rho\sigma\rho(i)}$. Noting that $\gamma_{\sigma\rho(i)}\gamma_{\sigma\rho\sigma\rho(i)}\gamma_{\sigma\rho(i)}^{-1}$ maps the edge $\gamma_{\sigma\rho(i)} e_{\sigma\rho\sigma\rho(i)}$ of $\gamma_{\sigma\rho(i)}\mathcal{F}$ to $\gamma_{\sigma\rho(i)} e_{\rho\sigma\rho(i)}$, we see that $\gamma_{\sigma\rho(i)}\gamma_{\sigma\rho\sigma\rho(i)}\mathcal{F}$ is the p-gon adjacent to $\gamma_{\sigma\rho(i)}\mathcal{F}$ sharing the edge $\gamma_{\sigma\rho(i)}\gamma_{\sigma\rho\sigma\rho(i)} e_{\sigma\rho\sigma\rho(i)} = \gamma_{\sigma\rho(i)} e_{\rho\sigma\rho(i)}$. Continuing in this manner anti-clockwise about the vertex v_i, we obtain

$$\gamma_{\sigma\rho(i)}\gamma_{\sigma\rho\sigma\rho(i)} \cdots \gamma_{(\sigma\rho)^q(i)}\mathcal{F} = \mathcal{F}$$

and

$$\gamma_{\sigma\rho(i)}\gamma_{\sigma\rho\sigma\rho(i)} \cdots \gamma_{(\sigma\rho)^q(i)} e_{(\sigma\rho)^q(i)} = e_i.$$

Since $\mathcal{F}$ was assumed to be a fundamental domain for Γ, the first equality shows that

$$\gamma_{\sigma\rho(i)}\gamma_{\sigma\rho\sigma\rho(i)} \cdots \gamma_{(\sigma\rho)^q(i)} = 1,$$

and thus the latter gives

$$(\sigma\rho)^q(i) = i.$$

This holds for any $i = 1, \ldots, p$, and hence $(\sigma\rho)^q = 1$.

Conversely, suppose we have an involution $\sigma \in S_p$ with $(\sigma\rho)^q = 1$. We generate the group Γ by using the edge-pairing isometries described by σ. Specifically, let γ_i be the unique orientation-preserving isometry of the hyperbolic plane which maps the edge e_i onto $e_{\sigma(i)}$ (and does not map $\mathcal{F}$ onto itself). Now let Γ be the subgroup of the full group of symmetries of the tessellation of type $\{p, q\}$ which is generated by $\gamma_1, \ldots, \gamma_p$. Note that we immediately have the relation $\gamma_i^{-1} = \gamma_{\sigma(i)}$. We need to show that Γ acts freely and transitively on the set of p-gons in the tessellation.

For any $\gamma \in \Gamma$, the p-gon adjacent to $\gamma\mathcal{F}$ and sharing the edge γe_i is $\gamma\gamma_{\sigma(i)}\mathcal{F}$. Furthermore, the generator $\gamma_{\sigma(i)}$ which appears in this context is clearly unique as such amongst the generators $\gamma_1, \ldots, \gamma_p$. This observation yields a one-to-one correspondence between words in these generators (not using inverses) and paths in the dual graph[1] of the tessellation beginning at $\mathcal{F}$. The correspondence associates the word $\gamma_{i_1} \dots \gamma_{i_n}$ to the path passing successively through the p-gons

---

[1] The dual graph of the tessellation is the graph whose vertex set is the set of p-gons in the tessellation, with two p-gons being joined by an edge if and only if the p-gons are adjacent.

$$\mathcal{F},\ \gamma_{i_1}\mathcal{F},\ \dots,\ \gamma_{i_1}\cdots\gamma_{i_n}\mathcal{F}.$$

Furthermore, two such paths are homotopic in the dual graph (relative to their endpoints) if and only if their corresponding words can be made equal using only the relations $\gamma_i \gamma_{\sigma(i)} = 1$.

The connectedness of the dual graph now immediately implies the transitivity of Γ acting on the set of p-gons of the tessellation. It remains to demonstrate the freeness of this action. In the context of the dual graph, it needs to be shown that every path beginning and ending at $\mathcal{F}$ corresponds to a word which multiplies in Γ to give the identity element.

Using the arguments from the first part of the proof, one can observe that the hypothesis $(\sigma\rho)^q = 1$ is precisely the condition that the word corresponding to the path around any given vertex v_i of $\mathcal{F}$ multiplies in Γ to give the identity. Let us call these "primitive paths". To complete the proof one only needs to observe that any path in the dual graph which begins and ends at $\mathcal{F}$ can be written as a concatenation of conjugates of these primitive paths.

□

Theorem 0.1 is now reduced to a combinatorial problem.

Lemma 0.3. *There exists an involution $\sigma \in S_p$ such that $(\sigma\rho)^q = 1$ if and only if q has a divisor $d \neq 1$ which is less than or equal to p.*

Proof. If $(\sigma\rho)^q = 1$ then q is a multiple of the order of $\sigma\rho$, which is greater than one and divides $p!$. This proves necessity.

For sufficiency, suppose $m \leq p$ is a divisor of q. We will explicitly construct an involution $\sigma \in S_p$ such that $\sigma\rho$ has order m. Put

$$p = am + r$$

for integers a and r with $0 \leq r < m$. Note that $a \geq 1$ since $m \leq p$. Now define σ as the following product of disjoint transpositions:

$$\sigma = \prod_{j=1}^{a-1} \Big(j(m-1)+1 \;\; p-(j-1) \Big) \prod_{k=0}^{r-1} \Big(p-a-2k \;\; p-a-(2k-1) \Big)$$

This formula should be interpreted using the convention that if $a = 1$ then the first product is empty, and if $r = 0$ then the second product is empty. Calculation reveals that the product $\sigma\rho$ decomposes as the following product of disjoint m-cycles:

$$\begin{aligned} \sigma\rho \quad = \quad & \prod_{j=1}^{a-1} \Big((j-1)(m-1)+1 \;\; (j-1)(m-1)+2 \;\; \dots \;\; j(m-1) \;\; p-(j-1) \Big) \cdot \\ & \Big((a-1)(m-1)+1 \;\; (a-1)(m-1)+2 \;\; \dots \\ & \qquad\qquad \dots \;\; p-a-2r \;\; p-a-(2r-1) \;\; p-a-(2r-3) \;\; \dots \\ & \qquad\qquad\qquad\qquad\qquad\qquad \dots \;\; p-a-1 \;\; p-a+1 \Big). \end{aligned}$$

□

The arguments above are not particular to hyperbolic geometry. The results hold equally in the Euclidean case ($1/p + 1/q = 1/2$) and the spherical case ($1/p + 1/q > 1/2$), where the Fuchsian group is replaced by a discrete group of orientation-preserving isometries of $\mathbb{R}^2$ or S^2, respectively. However, this does not add anything significant to the result since there are only finitely many such regular tessellations. In fact, one finds that all such regular tessellations of the plane and sphere are realisable as tessellations by fundamental domains of some discrete group of orientation preserving isometries with one exception — the icosahedron (type $\{3, 5\}$).

References

[1] S. Katok, "Fuchsian Groups", University of Chicago Press, 1992.

[2] H. Poincaré, Théorie des groupes Fuchsiens, Acta Math. 1 (1882) 1–62.